

Fig. B. 7.2t. Activities of selected radionuclides as a function of soil depth.

100 METERS



JANET

Fig. B.8.1.a.

100 METERS






Fig. B.8.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



100 METERS



Fig. B.8.1.f. Soil-sample locations.

100 METERS





Fig. B.8.1.g. Vegetation sample locations.

100 METERS



JANET

Fig. B.8.1.h. The gamma-ray exposure rates ( $\mu\text{R}/\text{hr}$ ) measured 1 m above the ground by the LiF thermoluminescent dosimeters (TLD). The numbers shown in parentheses denote the location identification numbers.

100 METERS




Fig. B.8.1.i. The average  $^{239}\text{Pu}$  activities (pCi/gm) in soil samples collected to a depth of 15 cm.

JANET

100 METERS




Fig. B.8.1.j. The average  $^{90}\text{Sr}$  activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.8.1.k.  $^{137}\text{Cs}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



100 METERS





Fig. B.8.1.m.  $^{60}\text{Co}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



100 METERS




Fig. B.8.1.o. Terrestrial animal sample locations.




Fig. B. 8. 2a. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2c. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2d. Activities of selected radionuclides as a function of soil depth.




Fig. B.8.2e. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2f. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2g. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2h. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2i. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2j. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2k. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.21. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2m. Activities of selected radionuclides as a function of soil depth.




Fig. B. 8.2n. Activities of selected radionuclides as a function of soil depth.

100 METERS



Fig. B.9.1.a.

100 METERS



Fig. B.9.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.9.1.d. The gamma background exposure rate ( $\mu\text{R}/\text{hr}$ ) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.9.1.f. Soil-sample locations.

100 METERS



Fig. B.9.1.g. Vegetation sample locations.

100 METERS




Fig. B.9.1.i. The average  $^{239}\text{Pu}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B.9.1.j. The average  $^{90}\text{Sr}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B.9.1.k.  $^{137}\text{Cs}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)



Fig. B.9.1.1. The average  $^{137}\text{Cs}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.9.1.m.  $^{60}\text{Co}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.9.1.n. The average  $^{60}\text{Co}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.



Fig. B.9.1.o. Terrestrial animal sample locations.




Fig. B. 9.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B. 9. 2b. Activities of selected radionuclides as a function of soil depth.




Fig. B.9.2c. Activities of selected radionuclides as a function of soil depth.



Fig. B.10.1.a.




Fig. B.10.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.10.1.d. The gamma background exposure rate ( $\mu\text{R}/\text{hr}$ ) at 1 m above the ground, measured with a portable NaI scintillation counter.




Fig. B.10.1.f. Soil-sample locations.

100 METERS




Fig. B.10.1.g. Vegetation sample locations.




Fig. B.10.1.1. The average  $^{239}\text{Pu}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS

[Scale bar]




Fig. B.10.1.j. The average  $^{89}\text{Sr}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.10.1.k.  $^{137}\text{Cs}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



Fig. B.10.1.1. The average  $^{137}\text{Cs}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.10.1.m.  $^{60}\text{Co}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.10.1.n. The average  $^{60}\text{Co}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.



Fig. B.10.1.o. Terrestrial animal sample locations.




Fig. B.10.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.10.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B.10.2c. Activities of selected radionuclides as a function of soil depth.




Fig. B.10.2d. Activities of selected radionuclides as a function of soil depth.




Fig. B.10.2e. Activities of selected radionuclides as a function of soil depth.

100 METERS




Fig. B.11.1.a.

PERCY




Fig. B.11.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.11.1.d. The gamma background exposure rate ( $\mu\text{R}/\text{hr}$ ) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.11.1.f. Soil-sample locations.




Fig. B.11.1.1. The average  $^{239}\text{Pu}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



3.6                    73                    47  
7.0                    67                    9.8

-N-




Fig. B.11.1.j. The average  $^{90}\text{Sr}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B.11.1.k.  $^{137}\text{Cs}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)




Fig. B.11.1.1. The average  $^{137}\text{Cs}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.11.1.m.  $^{60}\text{Co}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



0.08                    2.7                    2.9                    2.6  
0.39                                             0.23




Fig. B.11.1.n. The average  $^{60}\text{Co}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B.11.2a. Activities of selected radionuclides as a function of soil depth.

100 METERS



Fig. B.12.1.a.

100 METERS




Fig. B. 12.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.12.1.d. The gamma background exposure rate ( $\mu\text{R}/\text{hr}$ ) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.12.1.f. Soil-sample locations.




Fig. B.12.1.g. Vegetation sample locations.

100 METERS




Fig. B.12.1.1. The average  $^{239}\text{Pu}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.12.1.j. The average  $^{90}\text{Sr}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.12.1.k.  $^{137}\text{Cs}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)




Fig. B.12.1.1. The average  $^{137}\text{Cs}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.12.1.m.  $^{60}\text{Co}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)




Fig. B.12.1.n. The average  $^{60}\text{Co}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.



Fig. B.12.1.o. Terrestrial animal sample locations.




Fig. B.12.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.12.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B. 12. 2c. Activities of selected radionuclides as a function of soil depth.

100 METERS



Fig. B.13.1.a.

NANCY

100 METERS




Fig. B.13.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.13.1.d. The gamma background exposure rate ( $\mu\text{R}/\text{hr}$ ) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS



Fig. B.13.1.f. Soil-sample locations.

100 METERS




Fig. B.13.1.g. Vegetation sample locations.

100 METERS





Fig. B.13.1.1. The average  $^{239}\text{Pu}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



Fig. B.13.1.j. The average  $^{80}\text{Sr}$  activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.13.1.k.  $^{137}\text{Cs}$  isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)