

100 METERS



Fig. B.13.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.13.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.13.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B.13.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.13.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B.13.2c. Activities of selected radionuclides as a function of soil depth.




Fig. B.13.2d. Activities of selected radionuclides as a function of soil depth.

100 METERS



Fig. B.14.1.a.

OLIVE

100 METERS




Fig. B.14.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)




Fig. B.14.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.14.1.f. Soil-sample locations.

100 METERS




Fig. B.14.1.g. Vegetation sample locations.

100 METERS




Fig. B.14.1.1. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.14.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.14.1.k. ^{137}Cs isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.14.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.14.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.14.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.14.1.o. Terrestrial animal sample locations.




Fig. B.14.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.14.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B.14.2c. Activities of selected radionuclides as a function of soil depth.




Fig. B. 14.2d. Activities of selected radionuclides as a function of soil depth.

100 METERS



PEARL

Fig. B.15.1.a.

100 METERS



Fig. B.15.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.15.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS



Fig. B.15.1.f. Soil-sample locations.

100 METERS




Fig. B.15.1.g. Vegetation sample locations.

100 METERS




Fig. B.15.1.h. The gamma-ray exposure rates ($\mu\text{R/hr}$) measured 1 m above the ground by the LiF thermoluminescent dosimeters (TLD). The numbers shown in parentheses denote the location identification numbers.

100 METERS



Fig. B.15.1.1. The average ^{239}Pu activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS



Fig. B.15.1.j. The average ^{90}Sr activities (pCi/gm) in soil samples collected to a depth of 15 cm.




Fig. B.15.1.k. ^{137}Cs isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



Fig. B.15.1.1. The average ^{137}Cs activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.15.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.15.1.n. The average ^{60}Co activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.15.1.o. Terrestrial animal sample locations.




Fig. B.15.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B. 15. 2b. Activities of selected radionuclides as a function of soil depth.




Fig. B. 15.2c. Activities of selected radionuclides as a function of soil depth.




Fig. B.15.2d. Activities of selected radionuclides as a function of soil depth.




Fig. B. 15.2e. Activities of selected radionuclides as a function of soil depth.




Fig. B. 15. 2f. Activities of selected radionuclides as a function of soil depth.




Fig. B. 15.2g. Activities of selected radionuclides as a function of soil depth.




Fig. B.15.2h. Activities of selected radionuclides as a function of soil depth.

100 METERS



Fig. B.16.1.a.



Fig. B.16.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



Fig. B.16.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.



Fig. B.16.1.f. Soil-sample locations.

100 METERS



△△△ MESSERSCHMIDIA

Fig. B.16.1.g. Vegetation sample locations.

100 METERS

18 3.5 3.0 24 24

Fig. B.16.1.1. The average ^{238}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B.16.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



Fig. B.16.1.k. ^{137}Cs isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)



Fig. B.16.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.





Fig. B.16.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



0.37 0.29

16

0.82

15

Fig. B.16.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B.16. 2a. Activities of selected radionuclides as a function of soil depth.

100 METERS



Fig. B.17.1.a.

100 METERS




Fig. B.17.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.17.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.17.1.f. Soil-sample locations.

100 METERS



Fig. B.17.1.g. Vegetation sample locations.

100 METERS




Fig. B.17.1.1. The average ^{239}Pu activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS





Fig. B.17.1.j. The average ^{90}Sr activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.17.1.k. ^{137}Cs isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



Fig. B.17.1.1. The average ^{137}Cs activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.17.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.17.1.n. The average ^{60}Co activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.17.1.o. Terrestrial animal sample locations.




Fig. B.17.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.17.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B.17.2c. Activities of selected radionuclides as a function of soil depth.




Fig. B.17.2d. Activities of selected radionuclides as a function of soil depth.




Fig. B.17.2e. Activities of selected radionuclides as a function of soil depth.




Fig. B. 17.2f. Activities of selected radionuclides as a function of soil depth.




Fig. B.17.2g. Activities of selected radionuclides as a function of soil depth.




Fig. B.17.2h. Activities of selected radionuclides as a function of soil depth.




Fig. B.17.2i. Activities of selected radionuclides as a function of soil depth.




Fig. B.17.2j. Activities of selected radionuclides as a function of soil depth.

100 METERS



Fig. B.18.1.a.

ILDA

100 METERS




Fig. B.18.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



Fig. B.18.1.d. The gamma background exposure rate ($\mu\text{R}/\text{hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS



Fig. B.18.1.f. Soil-sample locations.

100 METERS




Fig. B.18.1.g. Vegetation sample locations.

100 METERS




Fig. B.18.1.1. The average ^{239}Pu activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.18.1.j. The average ^{90}Sr activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.18.1.k. ¹³⁷Cs isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



Fig. B.18.1.1. The average ^{137}Cs activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS



Fig. B.18.1.m. ^{60}Co Isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.18.1.n. The average ^{60}Co activities (pCi/gm) in soil samples collected to a depth of 15 cm.




Fig. B. 18. 2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.18.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B.18.2c. Activities of selected radionuclides as a function of soil depth.




Fig. B.18.2d. Activities of selected radionuclides as a function of soil depth.

100 METERS




Fig. B.19.1.a.

100 METERS




Fig. B.19.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.19.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.