

100 METERS



Fig. B.19.1.f. Soil-sample locations.

100 METERS




Fig. B.19.1.g. Vegetation sample locations.

100 METERS




Fig. B.19.1.1. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.19.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.19.1.k. ^{137}Cs isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.19.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.19.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



Fig. B.19.1.o. Terrestrial animal sample locations.




Fig. B.19.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.19.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B.19.2c. Activities of selected radionuclides as a function of soil depth.

100 METERS




Fig. B.20.1.a.

100 METERS




Fig. B.20.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.20.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.



Fig. B.20.1.f. Soil-sample locations.

100 METERS

△△△ MESSERSCHMIDIA

○○○ SCAEVOLA

◇ PISONIA

● COCONUT

◆ MORINDA

■ PANDANUS

VERA

Fig. B.20.1.g. Vegetation sample locations.

100 METERS



Fig. B.20.1.1. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.20.1.j. The average ^{80}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.20.1.k. ^{137}Cs isosexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.20.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.20.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.20.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B. 20.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.20.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B. 20. 2c. Activities of selected radionuclides as a function of soil depth.

100 METERS



Fig. B.21.1.a.

100 METERS



Fig. B.21.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.21.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS



Fig. B.21.1.f. Soil-sample locations.

100 METERS



Fig. B.21.1.g. Vegetation sample locations.

100 METERS




Fig. B.21.1.i. The average ^{239}Pu activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.21.1.j. The average ^{90}Sr activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.21.1.k. ¹³⁷Cs isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



Fig. B.21.1.1. The average ^{137}Cs activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.21.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.21.1.n. The average ^{60}Co activities (pCi/gm) in soil samples collected to a depth of 15 cm.




Fig. B.21.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.21.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B. 21. 2c. Activities of selected radionuclides as a function of soil depth.





Fig. B. 21. 2d. Activities of selected radionuclides as a function of soil depth.

100 METERS



YVONNE A

Fig. B.22.1.a.

100 METERS




Fig. B.22.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)


100 METERS



600 400 70 70 30 7
750 150

B.22.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS



B.22.1.f. Soil-sample locations.

100 METERS






Fig. B.22.1.g. Vegetation sample locations.

100 METERS




B.22.1.1.1. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 0 and 10 cm.

100 METERS





B.22.1.1.2. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 10 and 20 cm.

100 METERS



B.22.1.1.3. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 20 and 30 cm.

100 METERS



B.22.1.1.4. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 30 and 40 cm.

100 METERS



B.22.1.1.5. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 40 and 50 cm.

100 METERS




B.22.1.1.6. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 50 and 60 cm.

100 METERS




B.22.1.1.7. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 60 and 70 cm.

100 METERS



B.22.1.1.8. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 70 and 80 cm.

100 METERS




B.22.1.1.9. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 80 and 90 cm.

100 METERS




B22.1.1.10. The average ²³⁹Pu activities (pCi/gm) in soil samples collected between depths of 90 and 100 cm.

100 METERS



B.22.1.i.11. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 100 and 110 cm.

1.1.1.1



B.22.1.i.12. The average ^{239}Pu activities (pCi/gm) in soil samples collected between depths of 110 and 120 cm.

100 METERS




Fig. B.22.1.k. ^{137}Cs isoeposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.22.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.22.1.o. Terrestrial animal sample locations.




Fig. B. 22.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B. 22.2b. Activities of selected radionuclides as a function of soil depth.




Fig. B. 22.2c. Activities of selected radionuclides as a function of soil depth.




Fig. B. 22.2d. Activities of selected radionuclides as a function of soil depth.




Fig. B. 22.2e. Activities of selected radionuclides as a function of soil depth.




Fig. B.22.2f. Activities of selected radionuclides as a function of soil depth.




Fig. B. 22.2g. Activities of selected radionuclides as a function of soil depth.