

100 METERS




Fig. B.38.1.g. Vegetation sample locations.

100 METERS




Fig. B.38.1.1. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.38.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.38.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.38.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.38.1.o. Terrestrial animal sample locations.




Fig. B.38.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B. 38. 2b. Activities of selected radionuclides as a function of soil depth.




Fig. B. 38. 2c. Activities of selected radionuclides as a function of soil depth.




Fig. B.38.2d. Activities of selected radionuclides as a function of soil depth.




Fig. B.38.2e. Activities of selected radionuclides as a function of soil depth.

100 METERS




Fig. B.39.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

*Elevated levels may be due to nearby ^{60}Co source which was subsequently removed.

100 METERS




Fig. B.39.1.f. Soil-sample locations.




Fig. B.39.1.1. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.39.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.39.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



0.01

0.07

0.04

0.05 0.07

0.05

0.08

0.03 0.04 0.07
0.03 0.04 0.05
0.04 0.05 0.04
0.04 0.04 0.88
0.06 0.04
0.04

0.05

Fig. B.39.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.





Fig. B. 39. 2a. Activities of selected radionuclides as a function of soil depth.

100 METERS
[]



ELMER D

Fig. B.40.1.a.

100 METERS




Fig. B.39.1.a.

100 METERS




Fig. B.39.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS





Fig. B.40.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS





Fig. B.40.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS



B.40.1.f. Soil-sample locations.

100 METERS




Fig. B.40.1.1. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.40.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.40.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.40.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.



Fig. B.41.1.a.

100 METERS



Fig. B.41.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS



Fig. B.41.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS



Fig. B.41.1.f. Soil-sample locations.





Fig. B.41.1.g. Vegetation sample locations.



Fig. B.41.1.i. The average ^{239}Pu activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS



0.30

0.25

0.58

0.42

0.48

Fig. B.41.1.j. The average ^{90}Sr activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS



Fig. B.41.1.k. ^{137}Cs isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.41.1.1. The average ^{137}Cs activities (pCi/gm) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.41.1.m. ^{60}Co isoexposure and isoconcentration contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.41.1.n. The average ^{60}Co activities (pCi/gm) in soil samples collected to a depth of 15 cm.





Fig. B.41.2a. Activities of selected radionuclides as a function of soil depth.


100 METERS



FRED A

Fig. B.42.1.a.

100 METERS



FRED A

Fig. B.42.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.42.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.42.1.f. Soil-sample locations.

100 METERS




Fig. B.42.1.g. Vegetation sample locations.



Fig. B.42.1.i. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.42.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.42.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.42.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS
[] [] [] [] []




Fig. B.43.1.a.

100 METERS




Fig. B.43.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.43.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.43.1.f. Soil-sample locations.

100 METERS





Fig. B.43.1.i. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



1.5 0.60 0.26 0.56
 0.34 0.59 0.29 1.5

Fig. B.43.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.43.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.43.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.




Fig. B.43.2a. Activities of selected radionuclides as a function of soil depth.




Fig. B.43.2b. Activities of selected radionuclides as a function of soil depth.





Fig. B.43.2c. Activities of selected radionuclides as a function of soil depth.

100 METERS



FRED C

Fig. B.44.1.a.

100 METERS




Fig. B.44.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS





Fig. B.44.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground. measured with a portable NaI scintillation counter.


100 METERS



FRED C

Fig. B.44.1.f. Soil-sample locations.


100 METERS



0.05 0.07
0.07

Fig. B.44.1.i. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



1.5

0.60 0.26 0.56
0.34

FRED C

Fig. B.44.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.44.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS





Fig. B.44.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



0.03

0.08


0.02

0.05

0.04

Fig. B.44.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



FRED D

Fig. B.45.1.a.

100 METERS
|_|_|_|_|




Fig. B.45.1.b. Gross count isosexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.45.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.45.1.f. Soil-sample locations.

100 METERS



Fig. B.45.1.i. The average ^{239}Pu activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.45.1.j. The average ^{90}Sr activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.45.1.1. The average ^{137}Cs activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS



0.02



0.02

Fig. B.45.1.n. The average ^{60}Co activities (pCi/g) in soil samples collected to a depth of 15 cm.

100 METERS




Fig. B.46.1.a.

100 METERS




Fig. B.46.1.b. Gross count isoexposure contours. (Refer to alphabetic symbol key in this appendix.)

100 METERS




Fig. B.46.1.d. The gamma background exposure rate ($\mu\text{R/hr}$) at 1 m above the ground, measured with a portable NaI scintillation counter.

100 METERS




Fig. B.46.1.f. Soil-sample locations.